

LA-UR-02-7336

Approved for public release;  
distribution is unlimited.

*Title:* The anomalous suppression of  $\pi_2(1670) \rightarrow \pi b_1(1235) \pi$

*Author(s):* Philip R. Page, T-16

*Submitted to:* Proceedings of the XVIth International Conference  
on Particles and Nuclei  
Osaka, Japan, October 2002



## Los Alamos

NATIONAL LABORATORY

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

# The anomalous suppression of $\pi_2(1670) \rightarrow b_1(1235) \pi$

P.R. Page<sup>a\*</sup>

<sup>a</sup>Theoretical Division, MS B283, Los Alamos National Laboratory,  
Los Alamos, NM 87545, U.S.A.

We show that current experimental data indicate that the strong decay mode  $\pi_2 \rightarrow b_1 \pi$  is anomalously small (more than 3 times smaller than all other decay modes of the  $\pi_2$ ). This acts as a powerful discriminator for and against various decay models. Non-relativistic quark models with spin-1 pair creation, e.g.  $^3P_0$  (flux-tube breaking) and  $^3S_1$  and  $^3D_1$  (chromo-electric string-breaking) models, as well as lowest order one-boson (in this case  $\pi$ ) emission models, can accommodate the experimental data because of a quark-spin selection rule. Models that violate the selection rule, e.g. higher order one-boson emission decay mechanisms, as well as mixing with other Fock states and relativistic effects, may be constrained by the small  $\pi_2 \rightarrow b_1 \pi$  decay.

## 1. Experimental data on $\pi_2(1670) \rightarrow b_1(1235) \pi$

Recently, the VES Collaboration published for the first time an upper bound on the branching fraction  $\text{Br}[\pi_2 \rightarrow b_1 \pi] < 0.0019$  at the 97.7% confidence level. This branching fraction is measured in 37 GeV  $\pi^-$  collisions on a nucleus, in the reaction  $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$  [1]. This small branching fraction is consistent with preliminary data on the reaction  $\pi^- p \rightarrow \omega \pi^- \pi^0 p$  in 18 GeV  $\pi^-$  collisions on a proton from the E852 Collaboration [2].

The decay  $\pi_2 \rightarrow b_1 \pi$  is allowed by conservation of parity, angular momentum, charge conjugation, isospin and G-parity, and its strength should be compared with that of other decays which are allowed by the same quantum numbers, which are to an extraordinary degree conserved by the strong interactions. In order to show that the branching ratio is small for dynamical reasons, independent of any model, factors due to phase space and flavor should be removed. The standard expression of the partial width is [3]

$$\Gamma = \frac{p}{8\pi (2J_{\pi_2} + 1) m_{\pi_2}^2} |p^L f \mathcal{M}|^2 \quad (1)$$

where  $m_{\pi_2}$  and  $J_{\pi_2}$  are the mass and angular momentum of the decaying  $\pi_2$ , the decay momentum  $p$  is measured in the rest frame of the  $\pi_2$ ,  $L$  is the relative orbital angular momentum of the decay products, and  $p^L f \mathcal{M}$  is the decay amplitude. The amplitude without the phase space ( $p^L$ ) and flavor ( $f$ ) factors is labeled  $\mathcal{M}$ . In Table 1 we show the ratio of  $|\mathcal{M}|^2$  for the measured decay modes of the  $\pi_2$  to  $|\mathcal{M}|^2$  for the dominant decay mode ( $f_2 \pi$ ). A further refinement is to incorporate the usual suppression at large


---

\*This research is supported by the U.S. Department of Energy under contract W-7405-ENG-36.

Table 1

Partial widths with phase space and flavor removed relative to the dominant mode. Decay is assumed to happen in the bold-faced  $L$ -wave, since in all modes (except for  $f_2\pi$  where the D-wave is  $(0.18 \pm 0.06)^2 = (3.2 \pm 2.2)\%$  of the S-wave [3]) the contributions from the different partial waves are not known. Define the ratios  $R(X) = |\mathcal{M}(X)|^2 / |\mathcal{M}(f_2\pi)|^2$  and  $\tilde{R}(X) = |\tilde{\mathcal{M}}(X)|^2 / |\tilde{\mathcal{M}}(f_2\pi)|^2$ . The branching ratios do not add to unity, since ref. [3] only constrained some of the modes to add to unity. However, since  $R(X)$  and  $\tilde{R}(X)$  only measure relative branching ratios this is irrelevant to our conclusions.

Mode $X$	$p$ (GeV)	$L$	$f^2$	$Br(\pi_2 \rightarrow X)$ (%) [3]	$R(X)$	$\tilde{R}(X)$
$f_2\pi$	0.326	<b>S, D, G</b>	2	$56.2 \pm 3.2$	1.00	1.00
$\sigma\pi$	0.634	<b>D</b>	2	$13 \pm 6$	0.73	1.00
$\omega\rho$	0.308	<b>P, F</b>	2	$2.7 \pm 1.1$	0.53	0.53
$\rho(1450)\pi$	0.143	<b>P, F</b>	4	$< 0.36$	$< 0.36$	$< 0.33$
$\rho\pi$	0.649	<b>P, F</b>	4	$31 \pm 4$	0.33	0.46
$K\bar{K}^*$	0.450	<b>P, F</b>	2	$4.2 \pm 1.4$	0.27	0.30
$b_1\pi$	0.363	<b>D</b>	4	$< 0.19$	$< 0.09$	$< 0.09$

Figure 1. The results of Table 1 ( $R(X)$ ) plotted logarithmically.

momenta, i.e.  $\mathcal{M} = \exp(-\frac{1}{12}(\frac{p}{\beta})^2) \tilde{\mathcal{M}}$ , where  $\beta = 0.4$  GeV [4]. Table 1 also indicates these refined squared amplitude ratios. It is evident that the  $b_1\pi$  mode is suppressed by a factor of between 3 and 11 relative to the other modes, making it anomalously small. Since there is only an experimental upper bound on the  $b_1\pi$  mode, this suppression factor could be even larger, and we urge future experiments to put more restrictive bounds on this mode. An anomalously small mode will be distinctive on a logarithmic plot. As can be seen from Fig. 1, this is indeed the case.

## 2. Models that can accommodate $\pi_2(1670) \rightarrow b_1(1235)\pi$

The decay  $\pi_2 \rightarrow b_1\pi$  is particularly clean in the sense that it is only sensitive to OZI allowed decays (see Fig. 2). This is because OZI forbidden processes, which allow the creation of either the isovector  $\pi_2$ ,  $b_1$  or  $\pi$  out of only isoscalar gluons, are forbidden to the extent that isospin is conserved.

In non-relativistic quark-pair-creation models, where the OZI allowed decay process is modeled by an initial meson  $q\bar{q}'$  pair decaying to the two final meson pairs  $q\bar{Q}$  and  $Q\bar{q}'$ , a simple selection rule arises when all the mesons have quark-spin 0. If the pair creation of the  $Q\bar{Q}$  pair is with quark-spin  $S = 1$ , then conservation of quark-spin implies that the amplitude is zero. It can be argued that the decay  $\pi_2 \rightarrow b_1\pi$  is the only kinematically allowed decay involving experimentally discovered non-radially-excited mesons that can




Figure 2. The OZI allowed decay of an initial meson to two final mesons in various models.

test the selection rule.

It has been pointed out that a success of the non-relativistic flux-tube breaking model, where the created quark-antiquark pair has  ${}^3P_0$  quantum numbers (Fig. 2), i.e.  $S = 1$ , is the fact that  $\pi_2 \rightarrow b_1\pi$  is predicted to vanish due to the selection rule [5]. Other decay models where the created pair has  $S = 1$ , such as the non-relativistic chromo-electric string-breaking model where the pair has  ${}^3S_1$  or  ${}^3D_1$  quantum numbers (Fig. 2) [6], will also have this suppression. Both the  ${}^3P_0$  and  ${}^3S_1$  models involve a decay operator proportional to  $\vec{\sigma} \cdot \vec{p}$ , with  $\vec{\sigma}$  the Pauli matrices and  $\vec{p}$  a momentum operator.

While the flux-tube and chromo-electric string-breaking models have a  $q\bar{q}'$  pair decaying to the two final meson pairs  $q\bar{Q}$  and  $Q\bar{q}'$ , one of which is identified with the pseudoscalar boson, the one-pion emission model (Fig. 2) has either  $q \rightarrow Q\pi$  or  $\bar{q}' \rightarrow \bar{Q}'\pi$ . An expansion of the axial current in the lowest order one-boson coupling to the quark or antiquark gives a decay operator of the form  $\vec{\sigma} \cdot \vec{p}$  (Eqs. 2 and 28 or ref. [7]). This means that the boson is created from the quark with  $S = 1$ , so that the selection rule would also be valid for lowest order one-boson emission.

We conclude that the phenomenologically successful pair-creation model for light-light mesons (the  ${}^3P_0$  model) [5], the chromo-electric string-breaking model ( ${}^3S_1$  or  ${}^3D_1$  model), and the lowest order one-boson emission model, which has successfully been applied to the decay of heavy-light mesons [7,8], are consistent with the experimental decay width of  $\pi_2 \rightarrow b_1\pi$ .

### 3. Models possibly constrained by $\pi_2(1670) \rightarrow b_1(1235)\pi$

Higher order contributions in boson emission models do contain terms that are not of the form  $\vec{\sigma} \cdot \vec{p}$  which violate the selection rule. An example is interactions where *both* a pseudoscalar boson is emitted, *and* a particle is exchanged between the quark and antiquark in the initial meson (Eqs. 13, 38 and 39 or ref. [7]). The amplitudes corresponding to the higher order contributions can be similar in magnitude to those corresponding to the lowest order contribution (Table 4 of ref. [7]). We hence suggest that consistency with the small decay  $\pi_2 \rightarrow b_1\pi$  can be constraining for models which do not obey the selection rule, e.g. higher order contributions in one-boson emission models, and can provide a viability check on new decay mechanisms, e.g. the higher order contributions that were introduced [7] to cure problems with the lowest order contribution [7,8]. Although one-

gluon exchange (Fig. 2) violates the selection rule away from the non-relativistic limit,<sup>2</sup> it was found to be subdominant relative to the  $^3P_0$  model [9], so that it is not expected to be constrained by  $\pi_2 \rightarrow b_1\pi$ . If higher order contributions in boson emission models are inconsistent with  $\pi_2 \rightarrow b_1\pi$ , and one-gluon exchange is consistent, this could discriminate against the use of the boson exchange model.

#### 4. Further constraints due to $\pi_2(1670) \rightarrow b_1(1235)\pi$

The selection rule obtains when the  $\pi_2$ ,  $b_1$  and  $\pi$  are treated non-relativistically as quark-spin 0 mesons. In addition to decay models in the previous section, further breaking of the selection rule arises from:

- *Mixing with other Fock states:* The mixing of mesons participating in the decay with non-meson Fock states is constrained by the experimentally measured  $\pi_2 \rightarrow b_1\pi$  width. Examples of such mixing are: mixing between the quark-spin 0  $\pi_2$  meson and the quark-spin 1  $\pi_2$  hybrid meson expected nearby in mass, and non-mesonic Fock states in the Goldstone boson  $\pi$ .

- *Relativistic effects:* Relativistic corrections to the description of the mesons, which can introduce quark-spin 1 components in the  $\pi_2$ ,  $b_1$  and  $\pi$  mesons, are constrained by the experimentally measured  $\pi_2 \rightarrow b_1\pi$  width. We urge fully relativistic lattice QCD, QCD sum rule and Dyson-Schwinger Equation calculations of  $\pi_2 \rightarrow b_1\pi$ .

## REFERENCES

1. D.V. Amelin *et al.* (VES Collab.), Phys. At. Nucl. **62**, 445 (1999).
2. A. Popov (E852 Collab.), *Proc. of 9<sup>th</sup> Int. Conf. on Hadron Spectroscopy (HADRON 2001)* (Protvino, Russia, 25 Aug. – 1 Sep. 2001); M. Lu *et al.*, Bull. Am. Phys. Soc. **47**, 33 (2002).
3. K. Hagiwara *et al.* (Particle Data Group), Phys. Rev. D **66**, 010001 (2002).
4. See discussion in P.R. Page, Phys. Rev. D **60**, 057501 (1999).
5. T. Barnes, *Proc. of 3<sup>rd</sup> Workshop on Physics and Detectors for DAPHNE (DAPHNE 99)* (Frascati, Italy, 16-19 Nov 1999), eds. S. Bianco *et al.* (Frascati Physics Series Vol. 16, 1999), p. 503.
6. J.W. Alcock, M.J. Burfitt and W.N. Cottingham, Zeit. Phys. C **25** (1984) 161; P. Geiger and E.S. Swanson, Phys. Rev. D **50**, 6855 (1994).
7. K.O.E. Henriksson, T.A. Lähde, C.J. Nyfält and D.O. Riska, Nucl. Phys. A **686**, 355 (2001).
8. J.L. Goity and W. Roberts, Phys. Rev. D **60**, 034001 (1999).
9. E.S. Ackleh, T. Barnes and E.S. Swanson, Phys. Rev. D **54**, 6811 (1996).

---

<sup>2</sup>One-gluon exchange involves both a Coulomb and transverse interaction. The former has a simple  $\vec{\sigma} \cdot \vec{p}$  pair creation operator, but the latter involves *both*  $\vec{\sigma}$  pair creation and an additional term at the vertex where the quark or antiquark emits a gluon in Fig. 2 (Eqs. B5-B7 of ref. [9]). The additional term includes a  $\frac{1}{m} \vec{\sigma} \cdot \vec{p}$  contribution [9], which would mean that the overall transverse gluon interaction includes  $S = 1$  contributions at *both* interaction vertices of the gluon, giving rise to a violation of the selection rule.

# Los Alamos

NATIONAL LABORATORY

OK

## Technical Information Release

Group S-7, Mail Stop F674, Phone 7-5013, Fax 5-4251  
TA-3, SM-43, Room A115

2277

This form is to be completed and submitted to S-7, Classification Group, with copies as required of abstract or paper BEFORE you present or submit for publication any technical work of the Laboratory. All submittals to S-7 must also have the appropriate cover sheet.

**Allow three days for S-7 review.**

APPROVALS: Typed name and signature of responsible person required (Item 10).

### Required Copies

Cover sheet is required on all papers  
(Form 836 for LA-UR, Form 1756a-e for LA-CP)  
One of this form  
Two of unclassified abstract  
Three of unclassified paper  
One of classified or controlled abstract/paper

LA-UR/~~LA-CP~~

02-7336

1. Author(s) (full name and group affiliation)

Philip R. Page, 121963, T-16

2. Author(s) Signature and Date (optional for S-7)

3. Title of Article (in caps: spell out all symbols)

THE ANOMALOUS SUPPRESSION OF PI 2(1670) RIGHTARROW B1 (1235 PI

4. Type of Information

Abstract ☐ Full Paper ☒

Summary ☐ Poster ☐

Audio-Visual  
(abstract required) ☐

Vugraphs  
(abstract required) ☐

Other: ☐

Intended for: Journal ☐ Proceedings ☒ Meeting ☐ Talk ☐ Book ☐  
Book Chapter ☐ Electronic (e.g. e-print archive) ☐ Other: ☐

Particulars: (See instruction sheet #4.)

Proceedings of the XVIth International Conference on Particles and Nuclei, Osaka, Japan,  
October 2002

Controlled Meeting/Journal/Report: Yes ☐ No ☒

5. List the LA numbers of previous reports that might be useful to the reviewer

6. Research sponsored by:

DOE ☒ DOD ☐ NRC ☐ Other: ☐

Program Code used  
to fund this work: JAMP

DOE OSTI Distribution? Yes ☐ No ☐

7. Funding agency release required: Yes ☐ No ☒ Copy Attached: Yes ☐ No ☐

8. Deadline Date  
11/27/02

Telephone number for notification of release  
7-4835

Mail Stop  
B283

9. Derivative Classifier's Signature or DUSA Designator

Date

Classified ☐ Unclassified ☒ Unclassified, limited ☐ (Explain):

10. Typed/Printed Name of Responsible Party

Alan Bishop

Signature and Date

### Classification Group (S-7)

Date Received  
11-21-02

Subject or NRC Category  
72

Group/Author Notification of Release  
11/22/02 mjh kay.v.m.

Reviewed by

Date  
11/21/02

U ☒ C ☐ S ☐ RD ☐ FRD ☐ NSI ☐ UCN ☐ ECI ☐ OUO ☐  
Proprietary ☐ CRADA ☐ Guide \_\_\_\_\_ Duration \_\_\_\_\_

### Business and Patent Law

Patent Interest

Yes ☐ No ☒

Docket Number

Patent Law Reviewer and Date

Julian Spink 22 NOV 02

Comments: 4 pg's